

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

BC

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 460 607 A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 91109120.5

(22) Date of filing: 04.06.91

(61) Int. Cl.⁵: **C12P 21/08**, A61K 39/395,
G01N 33/577, G01N 33/574,
A61K 49/02, A61K 47/00,
C12N 15/12, C07K 15/14,
A61K 39/00

(30) Priority: 05.06.90 US 533371

(43) Date of publication of application:
11.12.91 Bulletin 91/50

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **BRISTOL-MYERS SQUIBB
COMPANY**
345 Park Avenue
New York, N.Y. 10154-0037(US)

(72) Inventor: Hellstrom, Ingegerd
3925 N.E. Surber Drive
Seattle, Washington 98105(US)

Inventor: Hellstrom, Karl Erik
3925 N.E. Surber Drive
Seattle, Washington 98105(US)
Inventor: Marquardt, Hans
9222 S.E. 46th Street
Mercer Island, Washington 98040(US)
Inventor: Johnston, Janet
9037 Palatine Avenue N.
Seattle, Washington 98103(US)

(74) Representative: Kinzebach, Werner, Dr. et al
Patentanwälte Reitsstötter, Kinzebach und
Partner Sternwartstrasse 4 Postfach 86 06 49
W-8000 München 86(DE)

(54) Novel monoclonal antibody to novel antigen associated with human tumors.

(57) The present invention is concerned with a novel monoclonal antibody which binds strongly to a protein antigen associated with human tumors, including carcinomas of the colon, breast, and lung, as well as melanomas. The antibody binds to normal human cells to a much lesser degree than to tumor cells. The antibody finds use both in diagnostic methods such as the detection of malignant cells associated with tumors and in therapeutic methods for treatment of humans with tumors. Also disclosed is a novel 70,000 - 75,000 dalton glycoprotein antigen found on the cell surface of human tumor cells. The amino terminal amino acid sequence of this antigen is:

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

in which X represents an unidentified amino acid.

EP 0 460 607 A2

The present invention relates to a novel monoclonal antibody and a novel antigen, and to methods for production and use of such novel monoclonal antibody reactive with human carcinoma cells. More specifically, the monoclonal antibody of this invention is reactive with the novel cell surface antigen, which is associated with a variety of human tumors including carcinomas of the breast, colon, as well as melanomas.

The monoclonal antibody of the present invention, designated L53, is reactive with a determinant of a glycoprotein antigen associated with non small cell lung carcinoma (NSCLC) cells and also with other carcinomas including breast and colon carcinomas, and melanomas. The L53 monoclonal antibody is suitable for both *in vivo* and *in vitro* clinical diagnostic purposes, such as the detection of malignant carcinomas. Additionally the antibody of the present invention is suited for therapeutic uses, for example to react with tumor cells, and in conjugates as a target-selective carrier of various agents which have anti-tumor effects including, but not limited to: chemotherapeutic drugs, toxins, immunological response modifiers, enzymes and radioisotopes. The antigen of the invention is also useful for therapeutic and diagnostic purposes.

Background of the Invention

Carcinomas cause millions of deaths annually. For example, lung carcinomas are responsible for the majority of deaths from cancer among men and are overtaking breast carcinomas as the most frequent cause of cancer death among women. Most cases of carcinomas are incurable by chemotherapy and radiation therapy unless radically removed in the early stages of the disease. There is thus a great need for methods of diagnosis and therapy of carcinomas of the breast, colon, ovary and lung, as well as for other malignant neoplasms such as melanomas.

Monoclonal antibodies reactive with carcinoma-associated antigens are known (see, e.g., Papsidero, *Semin. Surg. Oncol.*, 1 (4) 171-81 (1985); Schlom et al., *Important Adv. Oncol.*, 170-92 (1985); Allum et al., *Surg. Ann.*, 18:41-64 (1986); Houghton et al., *Semin. Oncol.* 13 (2):165-79 (1986); *Monoclonal Antibodies in Cancer: Advances for Diagnosis and Treatment*, Roth (ed.), Futura Publishing, Mt. Kisco, New York (1986); and *Cancer Diagnosis In Vitro Using Monoclonal Antibodies*, Kupchik (ed.) Marcel Dekker, Inc., New York, (1988)).

Most of the known monoclonal antibodies are reactive with several types of human carcinomas, while a few antibodies react with carcinomas derived from specific organs of the body, e.g., lung, breast, ovary, colon, stomach or pancreas. The target antigens are commonly glycoproteins or glycolipids (see, e.g., Hellstrom et al., *Cancer Research* 46:3917-23 (1986); and Fink et al., *Prog. Clin. Pathol.*, 9:121-33 (1984)). For example, monoclonal antibodies reactive with glycoprotein antigens on specific types of carcinomas include those described in United States Patent 4,737,579 (monoclonal antibodies to non-small cell lung carcinomas), United States Patent 4,753,894 (monoclonal antibodies to human breast cancer), United States Patent 4,579,827 (monoclonal antibodies to human gastrointestinal cancer), and United States Patent 4,713,352 (monoclonal antibodies to human renal carcinoma). Some monoclonal antibodies react with high molecular weight antigens which appear to be mucins. For example, monoclonal antibody B72.3 appears to recognize a tumor-associated oncofetal glycoprotein antigen of greater than 1,000 kd molecular weight that is selectively expressed on a number of different carcinomas. Thus, B72.3 has been shown to react with 84% of breast carcinomas, 94% of colon carcinomas, 100% of ovarian carcinomas and 96% of non-small-cell lung carcinomas (see Johnston, *Acta Cytol.*, 1 (5):537-56 (1987) and United States Patent 4,612,282, issued to Schlom et al.). Similarly, monoclonal antibody KC-4 recognizes an approximately 400-500 kd protein antigen expressed on a number of carcinomas, such as colon, prostate, lung and breast carcinoma (see United States Patent 4,708,930).

Monoclonal antibodies reactive with glycolipid antigens that are believed to be associated with certain tumor cells have also been disclosed. For example, Young et al., *J. Exp. Med.*, 150:1008-19 (1979) disclose the production of two monoclonal antibodies specific for asialo GM₂, a cell surface glycosphingolipid antigen that was established as a marker for BALB/c 3T3 cells transformed by Kirsten murine sarcoma virus. See, also, Kniep et al., *J. Immunol.*, 131 (3) 1591-94 (1983) and United States Patent 4,507,391 (monoclonal antibody to human melanoma).

In addition, monoclonal antibodies reactive with glycolipid antigens found on specific types of carcinoma cells include those described by Rosen et al., *Cancer Research*, 44:2052-61 (1984) (monoclonal antibodies to human small cell lung cancer); Varki et al., *Cancer Research*, 44:681-87 (1984) (monoclonal antibodies to human adenocarcinomas of the lung, stomach and colon and melanoma), and United States Patent 4,579,827 (monoclonal antibodies to human colon adenocarcinoma). See, also, Hellstrom et al., *Proc. Nat'l. Acad. Sci. USA*, 83:7059-63 (1986) which describes the L6 monoclonal antibody that recognizes a

carbohydrate antigen expressed on the surface of human non-small cell lung carcinomas, breast carcinomas and colon carcinomas.

Additional monoclonal antibodies exhibiting a reactivity to antigens found on a variety of tumor cells are greatly needed. This is because of the antigenic heterogeneity of most tumors which often necessitates, in diagnosis or therapy, the use of a combination of different monoclonal antibodies directed to the same tumor mass. Furthermore, monoclonal antibodies that display a high degree of reactivity with a wide range of tumors, while showing the absence of or only a very weak reactivity with normal tissues, are not common. Such antibodies would clearly be advantageous.

It is thus apparent that a monoclonal antibody reactive with an antigen expressed at high levels by a variety of tumors may become useful towards an earlier diagnosis of cancers, a better definition of the spread of the cancer, the immunological monitoring of cancer patients, as well as for development of improved methods for therapy of cancers. It is also apparent that monoclonal antibodies to novel cell surface molecules can be used for further definition of such molecules which may be of value for preparing immunogens in the form of cancer vaccines, and which may also have important cellular functions, for example, as receptors of hormones or growth factors or as molecules otherwise involved in intra- and intercellular communication. The antigens may even have enzymatic or growth factor activity by themselves.

Summary of the Invention

The present invention provides such a monoclonal antibody, L53, which is specific for a determinant site on a cell surface glycoprotein antigen, the L53 antigen, associated with a variety of human tumor cells, including lung, breast, and colon carcinoma and melanoma cells. Thus, the antibody of the invention can be useful for the diagnosis and therapy of tumors expressing the L53 antigen identified by antibody L53. The L53 antibody of the invention is of the IgG class, and IgG1 subclass and shows no significant reactivity with normal human cells.

The invention also comprises the novel L53 antigen identified by antibody L53 and the class of antibodies that bind, are immunospecific for or immunoreactive with this antigen. Further encompassed are methods for using the purified or cloned L53 antigen as a vaccine to immunize against certain carcinomas.

The antibody of the invention may be used in in vitro diagnostic methods for determining the presence of a malignant condition in human lung tissue and other human tissues. The methods involve examining the tissue for the presence of an antigen having the characteristics of the 70,000-75,000 dalton L53 glycoprotein reactive with antibody L53. For example, the tissue can be contacted with the L53 monoclonal antibody of the invention which defines a determinant site on a cell-associated antigen having the characteristics of the L53 antigen, or a functional equivalent or fragment of this antibody and any interactions of said antibody and antigenic determinants can be detected. One such method involves the determination of the presence of carcinoma cells in a specimen suspected of containing such cells. The specimen is contacted with the monoclonal antibody, which is capable of distinguishing such cells from other cell types which may be present in the specimen. The contact is carried out under conditions which promote the binding of the antibody to such cells. After contact, the presence or absence of binding of the antibody to the cells in the specimen is determined. This binding is related to the presence or absence of carcinoma cells in the specimen. Generally, the specimen is contacted with a labeled specific binding partner of the monoclonal antibody. This label is capable of producing a detectable signal. Alternatively, the monoclonal antibody itself may be labeled.

Another diagnostic method involves the in vivo localization of a tumor by administering to a patient a purified antibody or antibody fragment of the present invention labeled with an agent which gives a detectable signal. The localization is then detected using external scintigraphy, emission tomography or radionuclear scanning. This method can also provide better ways to stage cancer patients with respect to the extent of disease and to monitor changes in response to therapy.

The invention also has therapeutic applications, since the L53 antibody and similar antibodies can react with the L53 antigen that is expressed in high concentrations at the tumor cell surface. The monoclonal antibody of the invention may be used to prepare a composition for treating tumors. The composition comprises a therapeutically effective amount of the antibody in association with a pharmaceutically acceptable parenteral vehicle. The antibody of the invention can also be used in immunoconjugates as a carrier of various agents which have an antitumor effect, including, but not restricted to, chemotherapeutic drugs, toxins, immunological response modifiers, enzymes and radioisotopes.

Furthermore, the L53 antibody may be modified so that it can mediate antibody dependent cellular cytotoxicity (ADCC), that is, the antibody can kill carcinoma cells in the presence of human lymphocytes or

macrophages or it becomes cytolytic to tumor cells in the presence of human complement. Such modification can be accomplished, for example, by techniques recently developed for the production of chimeric antibodies as described by Oi et al., in *Biotechnologies* 4(3):214-221 (1986); and Fell et al., *Proc. Natl. Acad. Sci. USA* 86:8507-8511 (1989). Accordingly, genes coding for the variable region of the L53 antibody molecule are spliced together with human genes coding for the Fc region of an antibody with appropriate biological activity (such as the ability to activate human complement and mediate ADCC). Novel antibodies of mouse or human origin, can also be made to the L53 antigen having the appropriate biological functions.

The invention also comprises the novel L53 antigen characterized by a molecular weight of about 70,000-75,000 daltons and having an amino terminal amino acid sequence:

1	5	10	15	20	25	30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-M-X						

in which X represents an unidentified amino acid, and equivalents, identified by antibody L53 and the class of antibodies that bind to this antigen.

The invention includes methods for using the purified or cloned L53 antigen as a vaccine to immunize against certain tumors.

Detailed Description of Invention

In order that the invention herein described may be more fully understood, the following detailed description is set forth.

The present invention concerns a novel monoclonal antibody, designated L53, which is specifically reactive with an antigen (L53 antigen) localized at the membrane on human tumor cells including carcinomas of the lung, colon and breast, and melanoma cells, methods for producing the L53 monoclonal antibody and diagnostic and therapeutic methods employing the antibody. The L53 antibody reacts with a range of tumors while showing essentially no reactivity with normal human tissues or other types of tumors such as lymphomas.

The invention further concerns a novel cell surface glycoprotein antigen, designated L53 antigen, associated with human tumors of the lung, breast and colon, and melanomas and methods for using the L53 antigen.

The Monoclonal antibody L53 can be prepared by hybridoma fusion techniques or by techniques that utilize EBV-immortalisation technologies.

Hybridoma fusion techniques were first introduced by Kohler and Milstein (see, Kohler and Milstein, *Nature*, 256:495-97 (1975); Brown et al., *J. Immunol.*, 127 (2):539-46 (1981); Brown et al., *J. Biol. Chem.*, 255:4980-83 (1980); Yeh et al., *Proc. Nat'l. Acad. Sci. (USA)*, 76 (6):2927-31 (1976); and Yeh et al., *Int. J. Cancer*, 29:269-75 (1982)).

These techniques involve the injection of an immunogen (e.g., purified antigen or cells or cellular extracts carrying the antigen) into an animal (e.g., a mouse) so as to elicit a desired immune response (i.e., production of antibodies) in that animal. For example, human lung carcinoma cells from pleural effusions, cultured cells from explanted human non-small cell lung carcinomas (NSCLC), or cells from a normal fetal lung or lysates from such cells may be used as the immunogen. In the illustrative example herein, explanted cells from a NSCLC (human lung adenocarcinoma), lines CH3, 2981, 2707 and 2964 are used as the immunogen. The cells are injected repeatedly, for example, into a mouse and, after a sufficient time, the mouse is sacrificed and somatic antibody-producing lymphocytes are obtained. Antibody-producing cells may be derived from the lymph nodes, spleens and peripheral blood of primed animals. Spleen cells are preferred. Mouse lymphocytes give a higher percentage of stable fusions with the mouse myelomas described below. The use of rat, rabbit and frog somatic cells is also possible. The spleen cell chromosomes encoding desired immunoglobulins are immortalized by fusing the spleen cells with myeloma cells, generally in the presence of a fusing agent such as polyethylene glycol (PEG). Any of a number of myeloma cell lines may be used as a fusion partner according to standard techniques; for example, the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from the American Type Culture Collection (ATCC), Rockville, Maryland.

The resulting cells, which include the desired hybridomas, are then grown in a selective medium, such as HAT medium, in which unfused parental myeloma or lymphocyte cells eventually die. Only the hybridoma cells survive and can be grown under limiting dilution conditions to obtain isolated clones. The

supernatants of the hybridomas are screened for the presence of antibody of the desired specificity, e.g., by immunoassay techniques using the antigen that has been used for immunization. Positive clones can then be subcloned under limiting dilution conditions and the monoclonal antibody produced can be isolated. Various conventional methods exist for isolation and purification of the monoclonal antibodies so as to free them from other proteins and other contaminants. Commonly used methods for purifying monoclonal antibodies include ammonium sulfate precipitation, ion exchange chromatography, and affinity chromatography (see, e.g., Zola et al., in *Monoclonal Hybridoma Antibodies: Techniques And Applications*, Hurell (ed.) pp. 51-52 (CRC Press 1982)). Hybridomas produced according to these methods can be propagated in vitro or in vivo (in ascites fluid) using techniques known in the art (see, generally, Fink et al., *supra*, at page 123, Fig. 6-1).

Generally, the individual cell line may be propagated in vitro, for example in laboratory culture vessels, and the culture medium containing high concentrations of a single specific monoclonal antibody can be harvested by decantation, filtration or centrifugation. Alternatively, the yield of monoclonal antibody can be enhanced by injecting a sample of the hybridoma into a histocompatible animal of the type used to provide the somatic and myeloma cells for the original fusion. Tumors secreting the specific monoclonal antibody produced by the fused cell hybrid develop in the injected animal. The body fluids of the animal, such as ascites fluid or serum, provide monoclonal antibodies in high concentrations. As discussed by Cole et al., *supra*, when human hybridomas or EBV-hybridomas are used, it is necessary to avoid rejection of the xenograft injected into animals such as mice. Immunodeficient or nude mice may be used or the hybridoma may be passaged first into irradiated nude mice as a solid subcutaneous tumor, cultured in vitro and then injected intraperitoneally into pristane primed, irradiated nude mice which develop ascites tumors secreting large amounts of specific human monoclonal antibodies (see Cole et al., *supra*).

For certain therapeutic applications chimeric (mouse-human) or human monoclonal antibodies may be preferable to murine antibodies, because patients treated with mouse antibodies generate human antimouse antibodies. (Shawler et al., *J. Immunol.* 135:1530-35 (1985)). Chimeric mouse-human monoclonal antibodies reactive with the L53 antigen can be produced, for example, by techniques recently developed for the production of chimeric antibodies (Oi et al., *Biotechnologies* 4(3):214-221 (1986); Liu et al., *Proc. Nat'l. Acad. Sci. (USA)* 84:3439-43 (1987)). Accordingly, genes coding for the constant regions of the murine L53 antibody molecule are substituted with human genes coding for the constant regions of an antibody with appropriate biological activity (such as the ability to activate human complement and mediate ADCC). Novel antibodies of mouse or human origin, can be also made to the L53 antigen having the appropriate biological functions. For example, human monoclonal antibodies may be made by using the antigen, e.g. the L53 antigen of the invention, to sensitize human lymphocytes to the antigen in vitro followed by EBV-transformation or hybridization of the antigen-sensitized lymphocytes with mouse or human lymphocytes, as described by Borrebaeck et al. (*Proc. Nat'l. Acad. Sci. (USA)* 85:3995-99 (1988)).

According to a preferred embodiment, the antibody of this invention, designated L53, was produced via hybridoma techniques using four lung adenocarcinoma cell lines: CH3, 2981, 2707 and 2964, as the immunogen as described in the Example, *infra*. The L53 hybridoma, producing the L53 antibody, has been deposited with the ATCC, Rockville, Maryland, and has there been identified as follows:

L53 Accession No.: HB 10348

The L53 antibody is of the IgG1 subclass. The antibody displays a very strong reactivity with tumor cells of a variety of types, for example, carcinomas of the breast, lung, and colon, as well as with malignant melanoma. The L53 antibody shows no detectable binding to the T lymphoma cell line, CEM, and the B cell lymphoma line P3HR-1, and its binding to normal cells (epithelial, fibroblasts, endothelial, etc.) is weak or non-existent.

Thus, this antibody is superior to most known antitumor antibodies in its specificity for a range of tumor cells and in its high degree of specificity for tumor cells as compared to normal cells (see, e.g., Hellstrom et al., *Covalently Modified Antigens And Antibodies In Diagnosis And Therapy*, Quash/Rodwell (eds.), pp. 24-28 (Marcel Dekker, Inc., (1989); and Bagshawe, *Br. J. Cancer*, 48:167-75 (1983)).

It should be understood that the present invention encompasses the L53 antibody described above and any fragments thereof containing the active binding region of the antibody, such as Fab, F(ab')₂ and Fv fragments. Such fragments can be produced from the L53 antibody using techniques well established in the art (see, e.g., Rousseaux et al., in *Methods Enzymol.*, 121:663-69 Academic Press, (1986)).

In addition, the present invention encompasses antibodies that are capable of binding to the same antigenic determinant as the L53 antibody and competing with the L53 antibody for binding at that site. These include antibodies having the same antigenic specificity as the L53 antibody but differing in species origin, isotype, binding affinity or biological functions (e.g., cytotoxicity). For example, class, isotype and other variants of the antibody of the invention may be constructed using recombinant class-switching and

5

10

15

20

The amino terminal amino acid sequence of the novel L53 glycoprotein antigen is as follows:

26

30

35

40

45

5

al., *J. Immunol. Methods*, 42:11 (1981) and Allum et al., *supra*, at pp. 48-51). These assays, using the L53 antibody disclosed herein, can therefore be used for the detection in biological fluids of the L53 antigen with which the L53 antibody reacts and thus the detection of various carcinomas and melanomas in human patients. Thus, it is apparent from the foregoing that the L53 antibody of the invention can be used in most assays involving antigen-antibody reactions. These assays include, but are not limited to, standard RIA techniques, both liquid and solid phase, as well as ELISA assays, immunofluorescence techniques, and other immunocytochemical assays (see, e.g., Sikora et al. (eds.), *Monoclonal Antibodies*, pp. 32-52, Blackwell Scientific Publications, (1984)).

The L53 antibody of the invention is also useful for in vivo diagnostic applications for the detection of human tumors. One such approach involves the detection of tumors in vivo by tumor imaging techniques using the antibody labeled with an appropriate imaging reagent that produces detectable signal. Imaging reagents and procedures for labeling antibodies with such reagents are well known (see, e.g., Wensel and Meares, *Radio Immunoimaging and Radioimmunotherapy*, Esvier, New York (1983); Colcher et al., *Meth. Enzymol.*, 121:802-16 (1986)). The labeled antibody may be detected by a technique such as radionuclear scanning (see, e.g., Bradwell et al. in *Monoclonal Antibodies for Cancer Detection and Therapy*, Baldwin et al. (eds.), pp. 65-85, Academic Press (1985)).

L53 antibody can also be combined with other antibodies reactive with carcinomas such as the L6 antibody (Hellstrom et al., *Proc. Nat'l. Acad. Sci. USA* 83:7059-63 (1986)), to permit in vivo detection of non-small cell lung carcinomas such as large cell undifferentiated lung carcinoma, adenocarcinoma and squamous carcinoma.

Therapeutic Applications

The L53 antibody of the invention has a number of in vivo therapeutic applications. In addition to being used alone to target tumor cells, the antibody can be used in conjunction with an appropriate therapeutic agent to treat human cancer. For example, the antibody can be conjugated or linked to a therapeutic drug or toxin for delivery of the therapeutic agent to the site of the cancer. Techniques for conjugating such therapeutic agents to antibodies are well known (see, e.g., Arnon et al., *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56, Alan R. Liss, Inc., (1985); Hellstrom et al., in *Controlled Drug Delivery* (2nd ed.), Robinson et al. (eds.), pp. 623-53, Marcel Dekker, Inc., (1987); Thorpe, *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); and Thorpe et al., *Immunol. Rev.*, 62:119-58 (1982)). Since the L53 antibody is not easily internalized when cells are exposed to it in vitro, it may be preferable to target chemotherapeutic drugs to the tumor cells by coupling the antibody with an enzyme, e.g., using recombinant DNA techniques. When such conjugates are localized to the tumor, the enzyme can convert an inactive (nontoxic) prodrug which is administered after the conjugates have bound to the tumor cells, to an active anticancer drug. (See, e.g., Senter et al., *Proc. Nat'l. Acad. Sci. (USA)*, 85:4842-46 (1988)).

Alternatively, the antibody can be coupled to sources of high-energy radiation, e.g., a radioisotope such as ¹³¹I, which, when localized at the tumor site, results in a killing of several cell diameters (see, e.g., Order, in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 Academic Press, (1985)). According to yet another embodiment, the L53 can be conjugated to a second antibody to form an antibody heteroconjugate for the treatment of tumor cells as described by Segal in United States Patent 4,676,980.

Still other therapeutic applications for the L53 antibody of the invention include its use, either in the presence of complement or as part of an antibody-drug or antibody-toxin conjugate, to remove tumor cells from the bone marrow of cancer patients. According to this approach, autologous bone marrow may be purged ex vivo by treatment with the antibody and the marrow infused back into the patient (see, e.g., Ramsay et al., *J. Clin. Immunol.*, 8(2):81-88 (1988)).

Furthermore, chimeric or other recombinant L53 antibodies of the invention, as described earlier, may be used therapeutically, alone or as parts of various immunoconjugates. For example, a fusion protein comprising of at least the antigen-binding region of the L53 antibody joined to at least a functionally active portion of a second protein having anti-tumor activity, e.g., a lymphokine or tumor cell-inhibitor, may be used to treat human tumors in vivo. In addition, a chimeric L53 antibody wherein the antigen-binding region of L53 is joined to a human Fc region, e.g., IgG1, may be used to promote antibody-dependent cellular cytotoxicity or complement mediated cytotoxicity. Furthermore, recombinant techniques known in the art can be used to construct bispecific antibodies wherein one of the binding specificities of the antibody is that of L53 (see, e.g., United States Patent 4,474,893).

Finally, anti-idiotypic antibodies of the L53 antibody may be used therapeutically in active tumor

immunization and tumor therapy (see, e.g., Hellstrom et al., "Immunological Approaches To Tumor Therapy Monoclonal Antibodies, Tumor Vaccines, And Anti-Idiotypes", in Covalently Modified Antigens And Antibodies In Diagnosis And Therapy, supra, at pp. 35-41).

It is apparent therefore that the present invention encompasses pharmaceutical compositions, combinations and methods for treating human tumors. For example, the invention includes pharmaceutical compositions for use in the treatment of human tumors comprising a pharmaceutically effective amount of a L53 antibody and a pharmaceutically acceptable carrier. The compositions may contain the L53 antibody, either unmodified, conjugated to a therapeutic agent (e.g., drug, toxin, enzyme or second antibody) or in a recombinant form (e.g., chimeric or bispecific L53). The compositions may additionally include other antibodies or conjugates for treating carcinomas (e.g., an antibody cocktail).

The antibody compositions of the invention can be administered using conventional modes of administration including, but not limited to, intravenous, intraperitoneal, oral, intralymphatic or administration directly into the tumor. Intravenous administration is preferred.

The antibody compositions of the invention may be in a variety of dosage forms which include, but are not limited to, liquid solutions or suspensions, tablets, pills, powders, suppositories, polymeric microcapsules or microvesicles, liposomes, and injectable or infusible solutions. The preferred form depends upon the mode of administration and the therapeutic application.

The antibody compositions also preferably include conventional pharmaceutically acceptable carriers and adjuvants known in the art such as human serum albumin, ion exchangers, alumina, lecithin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, and salts or electrolytes such as protamine sulfate.

The most effective mode of administration and dosage regimen for the compositions of this invention depends upon the severity and course of the disease, the patient's health and response to treatment and the judgment of the treating physician. Accordingly, the dosages of the compositions should be titrated to the individual patient. Nevertheless, an effective dose of the antibody compositions of this invention may be in the range of from about 1 to about 5000 mg/m².

The novel antigen of the present invention, referred to as antigen L53 may also be used for therapeutic applications. The antigen can be purified from tumors or produced by recombinant DNA technology (Brown et al., copending U.S. Patent Application Serial No. _____, Attorney Docket No. 5624-123, filed on March 22, 1990, incorporated by reference herein). The gene coding for the L53 antigen may be cloned by methods which first enrich the mRNA of the L53 antigen. By one such method, polysomes (consisting of mRNA ribosomes and nascent polypeptide chains) can be purified by immunoaffinity chromatography with antibody that recognizes the L53 antigenic determinant on the nascent chain. The mRNA is isolated by immunoprecipitation with, e.g., L53 antibody and the cDNA is cloned in an appropriate expression vector. Alternatively, L53 antibody or antiserum to L53 antigen might be used to screen a cDNA library using an expression vector. The purified or cloned L53 antigen may be administered alone as an immunogen or together with a proper immunological adjuvant.

Purified or cloned L53 antigen may be used in the methods of the invention as a vaccine to immunize against certain tumors. Procedures for preparing such vaccines are known in the art (see, e.g., Estin et al., Proc. Nat'l. Acad. Sci. (USA), 85:1052 (1988)). Briefly, recombinant viruses are constructed for expression of the cloned tumor-associated antigen, for example L53 antigen. Cells infected with the recombinant viruses will express the tumor antigen at the surface of the cells together with the host's incompatibility antigens and immunogenic viral proteins. This favors the induction of cellular immunity which plays a key role in tumor rejection. A suitable virus, for example vaccinia virus derived from a plaque-purified virus of the Wyeth smallpox vaccine (New York City Board of Health strain), is used to construct a recombinant virus containing the coding sequence of the L53 antigen under control of the vaccinia virus "7.5 K" promoter (Hu et al., J. Virol. 62:176-180 (1988)). The recombinant virus may then be administered intravenously as a vaccine to protect against cancer.

50 Diagnostic Kits

The invention also encompasses diagnostic kits for carrying out the methods disclosed above: In one embodiment, the diagnostic kit comprises (a) monoclonal antibody L53 and (b) a conjugate of a specific binding partner for L53 antibody and a label for detecting bound antibody. The reagents may also include ancillary agents such as buffering agents and protein stabilizing agents, e.g. polysaccharides and the like. The diagnostic kit may further comprise, where necessary, other components of the signal-producing system including agents for reducing background interference, control reagents, an apparatus for conducting a test, etc. In another embodiment, the diagnostic kit comprises a conjugate of a monoclonal antibody to

the invention and a label capable of producing a detectable signal. Ancillary agents as mentioned above may also be present.

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting the scope of this invention in any manner.

EXAMPLE I

Preparation of the L53 Monoclonal Antibody

The L53 monoclonal antibody of the invention was produced using hybridoma fusion techniques described previously by Yeh et al., *Proc. Nat'l Acad. Sci. (USA)* (1979), *supra*. Briefly, a three month-old BALB/c mouse was immunized using explanted cultured cells from four human adenocarcinomas of the lung, designated CH3, 2981, 2707, and 2964, as the immunogen. The mouse received four (4) intraperitoneal (i.p.) injections and approximately 10^7 cells for each immunization. Three days after the last immunization, the spleen was removed, and the spleen cells were suspended in culture medium. The spleen cells were then fused with P3-x63-Ag8.653 mouse myeloma cells (ATCC No. CRL 1580), using polyethylene glycol (PEG), and the cell suspension grown in microtiter wells in selective HAT medium as described by Yeh et al., *Proc. Nat'l. Acad. Sci. (USA)*, *supra*. The mixture was seeded to form low density cultures originating from single fused cells or clones.

The supernatants from these hybridoma cultures were then screened for direct binding activity on the lung cancer cell lines CH3 and 2981 and against short-term cultures of human fibroblasts using an ELISA assay similar to that described by Douillard et al., *Meth. Enzymol.*, 92:168-74 (1983). According to this assay, the antigen (with which the antibody being screened for is reactive) is immobilized on microtiter plates and then incubated with hybridoma supernatants. If a supernatant contains the desired antibody, the antibody will bind to the immobilized antigen and is detected by addition of an anti-immunoglobulin antibody-enzyme conjugate and a substrate for the enzyme which leads to a measurable change in optical density.

For this example, lung cancer cells or control fibroblast cells or peripheral blood leukocytes (PBLs) were dispensed into a 96-well tissue culture plate (Costar, Cambridge, MA) and incubated overnight in a humid 37°C incubator (5% CO_2). The cells were then fixed with 100 μl of freshly prepared 1.0% glutaraldehyde to a final well concentration of 0.5% and incubated for 15 min at room temperature, followed by washing three times with 1 X PBS. The cells were next blocked for 30 min with 5% BSA in PBS and washed again three times with PBS. The supernatants from the hybridoma cultures were then added at 100 μl /well, the wells incubated for 1 hr at room temperature, and the cells washed three times with PBS. Next, goat anti-mouse horseradish peroxidase (Zymed, CA) diluted in 0.1% BSA and PBS was added to a concentration of 100 μl /well. The reaction mixture was incubated for either 1 hr at room temperature or 30 min at 37°C and the cells were then washed three times with PBS. Orthophenylenediamine (OPD) was then added at 100 μl /well and the plates incubated in the dark at room temperature for 5-45 min. Antibody binding to the cells was detected by a color change in the wells that occurred within 10-20 min. The reaction was stopped by adding 100 μl /well H_2SO_4 and the absorbance read in a Dynatech (Alexandria, VA) Microelisa autoreader at 492 nm.

The wells still positive on immunizing cell lines and negative on PBLs were tested by immunohistology technologies on immunizing cell line pellets and normal kidney, liver, and spleen tissue sections.

It should be noted that this assay can be performed using intact cells or purified soluble antigen or cellular extracts as the immobilized antigen. When soluble antigen or cell extracts were used as antigen, the antigen was initially plated at 50 μl /well in PBS and the plates were incubated overnight at room temperature before beginning the assay. When using intact cells as antigen, they may be used fresh or after fixation. In either case, the cells were initially plated at 10^4 cells at 100 μl /well in culture medium and incubated overnight in a 37°C incubator (5% CO_2).

Hybridomas which produced antibodies binding to the lung cancer cell line and not to the normal tissues were thus selected, cloned, expanded *in vitro*, and further tested for antibody specificity. Those hybridomas which produced antibody reactive with human lung cancer were recloned, expanded, and injected into pristane-primed 3-month old BALB/c mice, where they grew as ascites tumors.

Following this procedure, hybridoma cell line L53 was obtained, cloned and injected into mice to develop as an ascites tumor. As disclosed above, the L53 hybridoma has been deposited with the ATCC. Antibody secreted into the ascites was purified on protein A-Sepharose (see, e.g., Ey et al., *Immunochimistry*, 15:429-436 (1978)) or by gel filtration on Sephacryl S-300. Purified L53 antibody was used

for further characterization.

EXAMPLE II

5 Characterization of The L53 Monoclonal Antibody Isotype Determination

To determine the class of immunoglobulin produced by the L53 hybridoma, the following techniques were utilized:

10 a) Ouchterlony immunodiffusion.

An aliquot of supernatant of the L53 hybridoma cells was placed into the center well Of a 2.5% agar plate. Monospecific rabbit anti-mouse Ig isotype antibodies (Southern Biotechnology, Birmingham, AL) were placed in the outer wells and the plate was incubated for 24 hr at room temperature. Precipitation lines were
15 then read.

b) ELISA isotyping.

Dynatech Immulon 96-well plates were coated with goat anti-mouse Ig antibodies at 1 µg/ml concentration, 50 µl/well in PBS and left covered overnight at 4° C. The plates were washed with PBS/Tween 20, 0.05% and blocked with medium 100 µl/well for 1 hr at room temperature. After washing the plates, supernatants from the L53 hybridoma were added and incubated at room temperature for 1 hr. After washing with PBS containing bovine serum albumin (BSA) plates were incubated at 37° C for 2 hr with monospecific rabbit anti-mouse Ig isotype antibodies coupled to peroxidase (Zymed). After washing, plates
20 were incubated with 1 mg/ml o-phenylenediamine and 0.03% H₂O₂ in 0.1 M citrate buffer, pH 4.5. Optical density at 630 nm was determined on a Dynatec ELISA plate reader.

Based on these procedures, it was determined that the L53 monoclonal antibody is of the IgG1 isotype.

Binding Characteristics of The L53 Monoclonal Antibody

The subcellular localization of antigen was determined by measuring antibody binding to cells before or after permeabilization with non-ionic detergent. Antibodies binding to the cell surface of intact cultured cells were identified by direct fluorescence using the fluorescence activated cell sorter (FACS) II, as described by Hellstrom et al., Cancer Research 46:3817-3923 (1986). Briefly, for binding analyses using a FACS cell
35 sorter, 1 x 10⁶ cultured cells were aliquoted in 15% fetal bovine serum (FBS) in IMDM media (Gibco, Grand Island, NY) to a total volume of 500 µl/tube. The cells were centrifuged for 1.5 min on a Serofuge and the supernatant removed. 100 µl of the L53 monoclonal antibody at 10 µg/ml was added to each tube, the contents of which was then mixed and incubated on ice for 30 min. The reaction mixture was washed three times with 500 µl of 15% FBS/IMDM by centrifugation for 1.5 min on the Serofuge (tubes were blotted after
40 the third wash). Then, 50 µl of optimized FITC-conjugated goat anti-mouse IgG antibody (Tago, Burlingame, CA) diluted 1:25 in 15% FBS/IMDM was added to each tube and the reaction mixture was mixed and incubated for 30 min. The wash step was then repeated and after blotting of the tubes, each pellet was resuspended in 200-500 µl of PBS. Each sample was run on a Coulter Epics C FACS and the mean fluorescence intensity (MFI) was determined. From the MFI, the linear fluorescent equivalent (LFE) was
45 determined. The LFE of each test sample divided by the LFE of a negative control gave a ratio between the brightness of cells stained by specific vs. control antibody (1.0 = no difference in fluorescence, 2.0 = fluorescence twice as bright, etc.). To measure non-specific binding, parallel incubations were performed with unlabelled antibody as a competitor. The binding data are shown in Table 1 below.

50

55

Table 1
Binding of L53 Antibody to Various Cell Lines

	<u>Cell Lines</u>	<u>L53 Antibody Binding Ratio</u>
5		
	Lung Carcinoma	2707
		13
		CH3
		13
10		2964
		13
		2981
		13
	Colon Carcinoma	3347
		9
15		RCA
		14
		HCT 116
		15
		"C" line
		6
	Breast Carcinoma	3477
20		4
		3464
		5
		3396
		6
		CEM T lymphocytes
		1
25		P34R-1 B lymphocytes
		1

As Table 1 demonstrates, the L53 monoclonal antibody reacted with lung, breast and colon carcinoma cell lines, but did not react with T or B lymphoma lines nor with normal peripheral blood leukocytes.

Immunohistology

The PAP technique of L. A. Sternberger as described in *Immunochemistry*, pp. 104-69, John Wiley & Sons, New York (1979), as modified by Garrigues et al., *Int. J. Cancer*, 29:511-15 (1982), was used for immunohistological studies on frozen tissue sections. The target tissues for these tests were obtained at surgery and frozen within 4 hr of removal using isopentane precooled in liquid nitrogen. Tissues were then stored in liquid nitrogen or at -70° C until used. Frozen sections were prepared, air-dried, treated with acetone and dried again (see Garrigues et al., *supra*). Sections to be used for histologic evaluation were stained with hematoxylin. To decrease non-specific background, sections were preincubated with normal human serum diluted 1/5 in PBS (see Garrigues et al., *supra*). Mouse antibodies, rabbit anti-mouse IgG, and mouse PAP were diluted in a solution of 10% normal human serum and 3% rabbit serum. Rabbit anti-mouse IgG (Sternberger-Meyer Immunochemicals, Inc., Jarettsville, MD) was used at a dilution of 1/50. Mouse peroxidase-antiperoxidase complexes (PAP, Sternberger-Meyer Immunochemicals, Inc.) containing 2 mg/ml of specifically purified PAP were used at a dilution of 1/80.

The staining procedure consisted of treating serial sections with either specific antibody, i.e., L53, or a control antibody for 2.5 hr, incubating the sections for 30 min at room temperature with rabbit anti-mouse IgG diluted 1/50 and then exposing the sections to mouse PAP complexes diluted 1/80 for 30 min at room temperature. After each treatment with antibody, the slides were washed twice in PBS.

The immunohistochemical reaction was developed by adding freshly prepared 0.5% 3,3'-diaminobenzidine tetrahydrochloride (Sigma Chemical Co., St. Louis, MO) and 0.01% H₂O₂ in 0.05 M Tris buffer, pH 7.6, for 8 min (see Hellstrom et al., *J. Immunol.*, 127:57-60 (1981)). Further exposure to a 1% OsO₄ solution in distilled water for 20 min intensified the stain. The sections were rinsed with water, dehydrated in alcohol, cleared in xylene, and mounted on slides. Parallel sections were stained with hematoxylin.

The slides were each evaluated under code and coded samples were checked by an independent investigator. Typical slides were photographed by using differential interference contrast optics (Zeiss-Nomarski). The degree of antibody staining was evaluated as 0 (no reactivity), + (a few weakly positive cells), ++ (at least one third of the cells positive), +++ (most cells positive), ++++ (all cells strongly positive). Because differences between + and 0 staining were less clear cut than between + and ++

staining, a staining graded as ++ or greater was considered "positive". Both neoplastic and stroma cells were observed in tumor samples. The staining recorded is that of the tumor cells because the stroma cells were not stained at all or were stained much more weakly than the tumor cells.

Table 2 below presents the immunohistological staining of various tumor and normal tissue specimens using the L53 monoclonal antibody. As the table clearly demonstrates, the L53 antibody reacts with a wide range of human tumor specimens, and shows no reactivity or very weak reactivity with any of the number of normal human tissues tested.

TABLE 2
Immunoperoxidase Staining of Tumors and Normal
Tissue Specimens with L53 Monoclonal Antibody

15	<u>Tissue Type</u>	<u>Antibody Binding</u> (Number of Positive Tumors/ Total Number of Tumors Tested)
	CA. COLON	18/18
	CA. LUNG	15/26
	CA. BREAST	7/14
20	CA. OVARIAN	0/2
	CA. GASTRIC	0/2
	MELANOMA	5/5
	SARCOMA	1/5
25	<u>Normal Tissues:</u>	
	SPLEEN	0/6
	KIDNEY	2/10 ¹
	LIVER	1/10
	HEART	0/1
	OVARY	0/1
30	ADRENAL	0/1
	TESTIS	0/2
	BREAST	9/9 ²
	TONSIL	0/2
	SKIN	0/8
35	LUNG	0/9
	COLON	0/7
	BRAIN	0/6
	THYROID	0/7
40	LYMPH NODES	0/4
	PANCREAS	0/2
	ESOPHAGUS	0/2
	STOMACH	0/2

¹The only positive cells were scattered epithelial cells
²The only positive cells were scattered epithelial cells
in the breast glands

EXAMPLE III

L53 Antigen Recognized By L53 Antibody

55 Purification

In order to characterize the antigen reactive with the L53 monoclonal antibody, L53 antigen was isolated from 2981 cells (Oncogen, Seattle, WA) and from RCA cells (obtained from Dr. M. Brattain, Baylor

University, Houston, TX) and partially purified by immunoaffinity chromatography. L53 antigen was purified to homogeneity by SDS-PAGE and recovered from SDS-polyacrylamide gels by electroelution or electroblotting onto membranes.

Following electrophoresis, the SDS-polyacrylamide gel (12% acrylamide) was stained with Coomassie Brilliant Blue (0.05% by weight in 10% acetic acid and 30% isopropanol) and destained in a solution of acetic acid (5%, V:V) and methanol (17%, V:V). The stained L53 antigen band ($M_r = 70-75,000$ daltons) was excised with a razor blade and subjected to electroelution with a ECU-040 Electroelutor/Concentrator (C.B.S. Scientific Co., San Diego, CA), as described (Hunkapiller, et al., *Methods in Enzymology*, 91:227-236(1983)).

L53 antigen was also recovered from SDS-polyacrylamide gels (12% acrylamide) by electroblotting onto Immobilon membrane (Millipore Corp., Bedford, MA) using Mini-Transblot Electrophoretic Transfer Cell (BioRad Laboratories, Richmond, CA), as described (Matsudaira, J. *Biol. Chem.* 261:10035-1003(1987)). The membrane was stained with Coomassie Brilliant Blue, destained, and the stained L53 antigen band ($M_r = 70-75,000$ daltons) was excised with a razor blade for subsequent amino-terminal sequence analysis.

Sequence Analysis

Automated Edman degradation was performed on four preparations with 11 pmol (from 2981 cells), 16 pmol (from RCA cells), 21 pmol (from 2981 cells), and 25 pmol (from 2981 cells) ea., of L53 antigen in a pulsed-liquid protein sequencer (Model 475A, Applied Biosystems, Inc., Foster City, CA). The phenylthiohydantoin amino acid derivatives were analyzed by reversed-phase high performance liquid chromatography (HPLC) using a model 120A on-line HPLC unit (Applied Biosystems, Inc.) with a PTH C18 column (2.1 x 220 mm, Applied Biosystems, Inc.) and a sodium acetate/tetrahydrofuran/acetonitrile gradient for elution.

The amino-terminal sequence of L53 antigen is as follows:

1 5 10 15 20 25 30
D V V V O A P T Q V P G F L G D S V T L P X Y L Q V P N H X

in which X represents an amino acid that has not been identified.

A comparison of the 30-residue L53 amino-terminal sequence against the LOSPRO (May 1989), SWISSPROT (May 1989), and DIF (May 1989) data bases did not reveal significant sequence homology with any other known sequence. A similar comparison in the NBRF-PIR (release 24, May 1990) data base indicated a 28 residue identity with human poliovirus receptor H20A precursor.

The antigen recognized by L53 antibody is a novel protein of approximately 70,000 - 75,000 daltons molecular weight.

Immunological Characterization Radioimmunoprecipitation

a. 2669 and 2981 cells (20×10^6 cells, ea.) were surface-iodinated by the lactoperoxidase method with ^{125}I -Nal (New England Nuclear, Boston, MA) as described (Vitetta, E.S., et al., 1971, *J. Exp. Med.* 134:242-264). Radiolabeled membrane proteins were extracted with 1 ml lysis buffer (phosphate-buffered saline, pH 7.2, containing 1 mM EDTA, 1% Triton X-100, 0.1% SDS, 1% sodium deoxycholate, PMSF (phenylmethanesulfonyl fluoride; 10 $\mu\text{g/ml}$), and TLCX (N-tosyl-L-phenylalanine chloromethylketone; 2.0 $\mu\text{g/ml}$), for 30 min at 4°C . The detergent extract was precleared by incubation with 50 μl of a 50% suspension of Protein A-Sepharose CL-4B (Sigma Chemical Co., St. Louis, MO) and 5 μl of rabbit anti-mouse IgG (Zymed Laboratories, Inc., S. San Francisco, CA) for 30 min at 4°C . L53 antigen was immunoprecipitated by incubating the precleared cell lysate with 5 μl of purified L53 antibody (2 mg/ml) and 30 μl of a 50% suspension of Protein A-Sepharose CL-4B, previously coupled to rabbit anti-mouse IgG, for 2 h at 4°C . The immunoprecipitate was washed 4 times with 10 mM Tris-HCl buffer, pH 8.3, containing 0.5 M NaCl and 1% NP-40, analyzed by SDS-PAGE under reducing and non-reducing conditions and exposed to X-OMAT-Xray film (Kodak) with the aid of Lightening Plus intensifier screens.

b. 2669, RCA, and 2964 cells (1×10^6 cells/ml, ea.) were metabolically labeled with ^{35}S -methionine (0.25 mCi methionine/ml, Amersham Corp., Arlington Hts., IL) in RPMI 1640 media (methionine-free, RPMI 1640 Select-Amine Kit, GIBCO, Grand Island, NY), supplemented with 5% dialyzed fetal bovine serum, for 18h at 37°C . The cell pellets were washed 3 times with phosphate-buffered saline, pH 7.2, and extracted with 1 ml lysis buffer. L53 antigen was immunoprecipitated, as described above, analyzed by SDS-PAGE

under reducing and non-reducing conditions, and visualized by fluorography after impregnating the gel with Amplify (Amersham Corp.).

L53 antibody specifically precipitated L53 antigen with a Mr = 70-75,000 in both ¹²⁵I-Nal and ³⁵S-methionine-labeled cells.

5 The data demonstrate that the antigenic determinant recognized by L53 monoclonal antibody is localized on a unique single-chain protein with Mr = 70-75,000 daltons and is expressed on 2981, RCA, 2669, and 2964 cells.

10 It is apparent that many modifications and variations of this invention as set forth above may be made without departing from the spirit and scope. The specific embodiments described are given by way of example only and the invention is limited only by the terms of the appended claims.

Claims

- 15 1. A monoclonal antibody produced by hybridoma cell line ATCC No. HB 10348, which antibody binds to a determinant site on a cell surface glycoprotein antigen of human tumor cells and functional equivalents, binding fragments and immunocomplexes of said antibody.
2. The monoclonal antibody of claim 1 wherein said tumor cells are carcinoma cells or melanoma cells.
- 20 3. The monoclonal antibody of claim 2 wherein said carcinoma cells are selected from the group consisting of lung, colon and breast carcinoma cells.
4. The monoclonal antibody of claim 1 conjugated to a label capable of producing a detectable signal.
- 25 5. The monoclonal antibody of claim 4 wherein the label is selected from the group consisting of a radionuclide, an enzyme, a fluorescent agent and a chromophore.
6. A monoclonal antibody produced by hybridoma cell line ATCC No. HB 10348, which antibody binds to a cell determinant site on a cell surface glycoprotein antigen of human tumor cells, said antigen
- 30 characterized by a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis, and having an amino terminal amino acid sequence as follows:

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

35

in which X represents an unidentified amino acid and functional equivalents, binding fragments and immunocomplexes of said antibody.

- 40 7. A monoclonal antibody produced by a hybridoma cell line formed by fusion of a myeloma cell and a cell capable of producing antibody which binds to a determinant on a cell surface glycoprotein antigen of human tumor cells, said antigen having a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis, and having an amino terminal amino acid sequence as follows:

45

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

50 in which X represents an unidentified amino acid and functional equivalents, binding fragments and immunocomplexes of said antibody.

8. The monoclonal antibody of claim 7 which is of class IgG; of subclass IgG1 or is a murine antibody.
- 55 9. A monoclonal antibody reactive with a determinant site on a cell surface glycoprotein antigen associated with human tumor cells, said antigen characterized by a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis and having an amino terminal amino acid sequence as follows:

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-M-X

5 in which X represents an unidentified amino acid, and functional equivalents, binding fragments and immunocomplexes of said antibody.

10. The monoclonal antibody of claim 9 consisting of the L53 monoclonal antibody produced by hybridoma cell line ATCC No. HB 10348.

10 11. The monoclonal antibody of claim 9 which is a human antibody or which is a mouse-human antibody.

12. A composition for treating tumors comprising a therapeutically effective amount of the antibody of anyone of claims 1 to 3 and 6 to 11, in particular of claim 6 or 9, in association with a pharmaceutically acceptable parenteral vehicle.

13. An immunoassay for the detection of human tumors comprising:
a) combining a monoclonal antibody reactive with a cell surface glycoprotein antigen associated with human tumor cells, said antigen characterized by a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis and having an amino terminal amino acid sequence as follows:

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-M-X

25 in which X represents an unidentified amino acid, with a sample of tumor cells said antibody labeled so as to be capable of detection; and
b) assaying for said labeled monoclonal antibody binding to tumor cells associated with said antigen.

30 14. The immunoassay of claim 13 wherein said monoclonal antibody is the L53 antibody produced by hybridoma cell line ATCC No. HB 10348.

15. The immunoassay of claim 13 wherein said antibody is labelled with a label selected from the group consisting of a radionuclide, an enzyme, a fluorescent agent and a chromophore.

16. A method for detecting tumors which comprises: contacting the monoclonal antibody of claim 1, 6, 7 or 9 with a human tissue or fluid sample, and detecting interaction of said antibody with any antigenically corresponding tumor cells or antigenic determinants thereof in said sample.

40 17. The method of claim 16 wherein said tumor cells are lung carcinoma cells and the human tissue is lung tissue or

wherein said tumor cells are breast carcinoma cells and the human tissue is breast tissue or

45 wherein said tumor cells are colon carcinoma cells and the human tissue is colon tissue or

wherein said tumor cells are melanoma cells.

50 18. The method according to claim 16, wherein the interaction of said monoclonal antibody with said tumor cells is detected by immunohistological staining.

19. A composition for localizing human tumors in vivo, obtainable by

55 a) purifying the monoclonal antibody of claim 1, 6, 7 or 9;
b) radiolabeling said antibody; and
c) incorporating said antibody in a suitable carrier.

20. A composition for the treatment of tumors, obtainable by

- a) purifying the monoclonal antibody of claim 1, 6, 7 or 9;
- b) conjugating said monoclonal antibody to a cytotoxic agent, a toxin, or a radiopharmaceutical; and
- c) incorporating said conjugated monoclonal antibody in a suitable carrier.

- 5 21. The method of claim 20 wherein said monoclonal antibody is an anti-idiotypic antibody.
22. A continuous cell line which produces a monoclonal antibody, said monoclonal antibody characterized by the capability of binding to a determinant site on a cell surface glycoprotein antigen of human carcinoma cells, which comprises: a hybridoma of a lymphocyte derived from a mouse immunized with carcinoma cells or an immunogenic determinant thereof and a mouse myeloma cell.
- 10 23. A continuous cell line which produces a monoclonal antibody, said monoclonal antibody characterized by the capability of binding to a determinant site on a cell surface glycoprotein antigen of human carcinoma cells, which comprises: a hybridoma of a lymphocyte derived from a human with carcinoma and a myeloma cell.
- 15 24. Hybridoma cell line ATCC No. HB 10348 which produces a monoclonal antibody capable of binding to a determinant on a cell surface glycoprotein antigen of human tumor cells.
- 20 25. Hybridoma cell line ATCC No. HB 10348 formed by fusing a P3-x63-Ag8.653 mouse myeloma cell with a mouse splenocyte obtained from a BALB/c mouse immunized with lung adenocarcinoma CH3, 2981, 2907 and 2964 cells which produces a monoclonal antibody which binds to a determinant of a cell surface glycoprotein antigen of human tumor cells having a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis and having an amino terminal amino acid sequence as follows:
- 25

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

- 30 in which X represents an unidentified amino acid.

26. A continuous cell line which produces a monoclonal antibody being characterized by the capability of binding to a determinant site on a cell surface glycoprotein antigen associated with tumor cells, said antigen having a molecular weight of about 70,000 to about 75,000 daltons having an amino terminal amino acid sequence as follows:
- 35

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

- 40 in which X represents an unidentified amino acid, which comprises: a hybridoma of a lymphocyte capable of producing antibody against said antigen and a myeloma cell.

- 45 27. A glycoprotein antigen in substantially purified form, said antigen derived from human tumor cells and having a molecular weight of about 70,000 to about 75,000 daltons as determined by polyacrylamide gel electrophoresis, and having an amino terminal amino acid sequence as follows:

1 5 10 15 20 25 30
D-V-V-V-Q-A-P-T-Q-V-P-G-F-L-G-D-S-V-T-L-P-X-Y-L-Q-V-P-N-H-X

- 50 in which X represents an unidentified amino acid and immunocomplexes of this antigen.

- 55 28. A vaccine for use in immunization against tumors comprising a recombinant virus including the DNA coding for the antigen of claim 27,

in particular wherein said virus is vaccinia virus.

29. A diagnostic kit comprising:

- a) the monoclonal antibody of claim 1, 6, 7 or 9; and
- b) a conjugate of a detectable label and a specific binding partner of the monoclonal antibody of (a) above.

5

30. The diagnostic kit of claim 29 wherein the label is selected from the group consisting of enzymes, radiolabels, chromophores and fluorescers.

10

15

20

25

30

35

40

45

50

55